

# Children's Mucinex® Cold, Cough & Sore Throat Liquid (Mixed Berry Flavor)

## Drug Facts

### Active ingredients (in each 10 mL) Purposes

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

### Uses

- temporarily relieves these common cold and flu symptoms:
  - nasal congestion
  - stuffy nose
  - cough due to minor throat and bronchial irritation
  - the intensity of coughing
  - the impulse to cough to help your child get to sleep
  - minor aches and pains
  - sore throat
  - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes
- more than 5 doses in 24 hours, which is the maximum daily amount
  - with other drugs containing acetaminophen
- Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

### Ask a doctor before use if your child has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not use more than directed (see Overdose warning)

## Drug Facts (continued)

### Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

### Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

### Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- shake well before using
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- children 6 to under 12 years of age: 10 mL in dosing cup provided
- children under 6 years of age: do not use

### Other information

- each 10 mL contains: sodium 6 mg
- tamper evident: do not use if neckband on bottle cap is broken or missing.
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

**Inactive ingredients** anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate,\* xanthan gum \*may contain this ingredient

### Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.