

Maximum Strength Mucinex® Fast-Max® Cold, Flu & Sore Throat Liquid Gels

Drug Facts

Active ingredients (in each liquid gel)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - headache
 - minor aches and pains
 - temporarily reduces fever
 - nasal congestion
 - cough
 - sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- heart disease
- high blood pressure

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and older: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol, titanium dioxide, water

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.