

Maximum Strength Mucinex® Fast-Max®

DAY TIME Severe Cold and NIGHT TIME Cold & Flu Caplets

Drug Facts

Active ingredients (in each caplet) Purposes
Mucinex FAST-MAX DAY TIME Severe Cold

Acetaminophen 325 mg.....Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg.....Cough suppressant
 Guaifenesin 200 mg.....Expectorant
 Phenylephrine HCl 5 mg.....Nasal decongestant

Active ingredients (in each caplet) Purposes
Mucinex FAST-MAX NIGHT TIME Cold & Flu

Acetaminophen 325 mg.....Pain reliever/fever reducer
 Diphenhydramine HCl 25 mg.....Antihistamine
 Phenylephrine HCl 5 mg.....Nasal decongestant

Uses ■ temporarily relieves these common cold and flu symptoms:

- nasal congestion
- cough (*Mucinex FAST-MAX DAY TIME Severe Cold only*)
- minor aches and pains ■ headache ■ sore throat
- runny nose and sneezing (*Mucinex FAST-MAX NIGHT TIME Cold & Flu only*)
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (*Mucinex FAST-MAX DAY TIME Severe Cold only*)
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs that contain acetaminophen
- 3 or more alcoholic drinks daily while using this product

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin (*Mucinex FAST-MAX NIGHT TIME Cold & Flu only*)
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease ■ heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma (*Mucinex FAST-MAX NIGHT TIME Cold & Flu only*)
- a breathing problem such as emphysema or chronic bronchitis (*Mucinex FAST-MAX NIGHT TIME Cold & Flu only*)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema (*Mucinex FAST-MAX DAY TIME Severe Cold only*)
- cough that occurs with too much phlegm (mucus) (*Mucinex FAST-MAX DAY TIME Severe Cold only*) ▶

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (*Mucinex FAST-MAX NIGHT TIME Cold & Flu only*)

When using this product

- **do not use more than directed**
- excitability may occur, especially in children (*Mucinex FAST-MAX NIGHT TIME Cold & Flu only*)
- marked drowsiness may occur (*Mucinex FAST-MAX NIGHT TIME Cold & Flu only*)
- alcohol, sedatives, and tranquilizers may increase drowsiness (*Mucinex FAST-MAX NIGHT TIME Cold & Flu only*)
- avoid alcoholic drinks (*Mucinex FAST-MAX NIGHT TIME Cold & Flu only*)
- be careful when driving a motor vehicle or operating machinery (*Mucinex FAST-MAX NIGHT TIME Cold & Flu only*)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be a sign of a serious condition. (*Mucinex FAST-MAX DAY TIME Severe Cold only*)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients (Mucinex FAST-MAX DAY TIME Severe Cold)

croscarmellose sodium, crospovidone, FD&C Red #40 aluminum lake, FD&C Yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol 3350, polyvinyl alcohol, povidone, talc, titanium dioxide

Inactive ingredients (Mucinex FAST-MAX NIGHT TIME Cold & Flu)

corn starch, croscarmellose sodium, crospovidone, FD&C Blue #1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, triacetin

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.