

Maximum Strength Mucinex® Fast-Max®

Day Time Severe Cold / Night Time Cold & Flu Caplets

Drug Facts

Active ingredients (in each caplet)	Purposes
Mucinex FAST-MAX DAY TIME Severe Cold	
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Active ingredients (in each caplet)	Purposes
Mucinex FAST-MAX NIGHT TIME Cold & Flu	
Acetaminophen 325 mg	Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg	Antihistamine/cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses	temporarily relieves these common cold and flu symptoms:
<ul style="list-style-type: none"> ■ minor aches and pains ■ nasal congestion ■ runny nose (NIGHT TIME only) ■ sneezing (NIGHT TIME only) 	<ul style="list-style-type: none"> ■ cough ■ headache ■ sore throat

Drug Facts (continued)

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (DAY TIME only)
- controls cough to help you get to sleep (NIGHT TIME only)
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ▶

Drug Facts (continued)

- with any other drug containing diphenhydramine, even one used on the skin (NIGHT TIME only)
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (NIGHT TIME only)
- breathing problems such as emphysema or chronic bronchitis (NIGHT TIME only)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (NIGHT TIME only) ▶

Drug Facts (continued)

When using this product

- do not use more than directed
- excitability may occur, especially in children (NIGHT TIME only)
- marked drowsiness may occur (NIGHT TIME only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (NIGHT TIME only)
- avoid alcoholic drinks (NIGHT TIME only)
- be careful when driving a motor vehicle or operating machinery (NIGHT TIME only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. ▶

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients (Mucinex FAST-MAX DAY TIME Severe Cold)

croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol 3350, polyvinyl alcohol, povidone, talc, titanium dioxide

Inactive ingredients (Mucinex FAST-MAX NIGHT TIME Cold & Flu)

croscarmellose sodium, crospovidone, FD&C blue #1, FD&C blue #2, methacrylic acid – ethyl acrylate copolymer (1:1) type A, mica, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone K29/32, sodium bicarbonate, talc, titanium dioxide ▶

Drug Facts (continued)

Questions?

1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.