

Maximum Strength Mucinex® Fast-Max® Day Time Severe Cold / Night Time Cold & Flu Liquid

Drug Facts

Active ingredients (in each 20 mL) Purposes Mucinex FAST-MAX DAY TIME Severe Cold

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Drug Facts (continued)

Active ingredients (in each 20 mL) Purposes Mucinex FAST-MAX NIGHT TIME Cold & Flu

Acetaminophen 650 mg	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg	Antihistamine/ cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Drug Facts (continued)

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - runny nose and sneezing (NIGHT TIME only)
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (DAY TIME only)
- controls cough to help you get to sleep (NIGHT TIME only)
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin (NIGHT TIME only)
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma (NIGHT TIME only)
- trouble urinating due to an enlarged prostate gland
- breathing problems such as emphysema or chronic bronchitis (NIGHT TIME only)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (NIGHT TIME only)

Drug Facts (continued)

When using this product ■ do not use more than directed
■ excitability may occur, especially in children (NIGHT TIME only)

- marked drowsiness may occur (NIGHT TIME only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (NIGHT TIME only)
- avoid alcoholic drinks (NIGHT TIME only)
- be careful when driving a motor vehicle or operating machinery (NIGHT TIME only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided with product
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- dosing cup provided
- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients (Mucinex FAST-MAX DAY TIME Severe Cold)

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum *may contain this ingredient

Inactive ingredients (Mucinex FAST-MAX NIGHT TIME Cold & Flu)

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum *may contain this ingredient