

Mucinex®

Drug Facts

Active ingredient (in each extended-release bi-layer tablet)	Purpose
---	----------------

Guaifenesin 600 mg.....	Expectorant
-------------------------	-------------

Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Do not use ■ for children under 12 years of age

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Stop use and ask a doctor if ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: 1 or 2 tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if carton is open or if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store between 20-25°C (68-77°F)

Inactive ingredients carbomer 934P, NF; FD&C blue #1 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF