

## Drug Facts

### Active ingredients (in each liquid gel) Purposes **Mucinex SINUS-MAX DAY**

Acetaminophen 325 mg.....Pain reliever  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Guaifenesin 200 mg.....Expectorant  
Phenylephrine HCl 5 mg.....Nasal decongestant

### Active ingredients (in each liquid gel) Purposes **Mucinex SINUS-MAX NIGHT**

Acetaminophen 325 mg.....Pain reliever  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Doxylamine succinate 6.25 mg.....Antihistamine  
Phenylephrine HCl 5 mg.....Nasal decongestant

### Uses

- temporarily relieves:
  - nasal congestion
  - headache
  - cough
  - minor aches and pains
  - sinus congestion and pressure
  - runny nose and sneezing (**NIGHT only**)
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (**DAY only**)

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (**NIGHT only**)

## Drug Facts (continued)

- breathing problems such as emphysema or chronic bronchitis (**NIGHT only**)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**NIGHT only**)

### When using this product

- do not use more than directed
- excitability may occur, especially in children (**NIGHT only**)
- marked drowsiness may occur (**NIGHT only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**NIGHT only**)
- avoid alcoholic drinks (**NIGHT only**)
- be careful when driving a motor vehicle or operating machinery (**NIGHT only**)

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and older: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

### Other information

- store between 20-25°C (68-77°F)
- avoid excessive heat

### Inactive ingredients (DAY only)

FD&C yellow no. 6, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol, titanium dioxide, water

### Inactive ingredients (NIGHT only)

D&C yellow no. 10, FD&C blue no. 1, gelatin, glycerin, lecithin (soy), mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol, titanium dioxide, water