

Maximum Strength Mucinex[®] Sinus-Max[®]

Severe Congestion Relief/ Night Time Congestion & Cough Liquids

Drug Facts

Active ingredients (in each 20 mL) Purposes Mucinex SINUS-MAX Day Time

Acetaminophen 650 mg	Pain reliever
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Active ingredients (in each 20 mL) Purposes Mucinex SINUS-MAX Night Time

Acetaminophen 650 mg	Pain reliever
Diphenhydramine HCl 25 mg	Antihistamine/ cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves:
 - headache
 - minor aches and pains
 - controls cough to help you get to sleep (**Night Time only**)
 - runny nose and sneezing (**Night Time only**)
 - temporarily promotes nasal and/or sinus drainage
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (**Day Time only**)
- nasal congestion
- cough (**Night Time only**)
- sinus congestion and pressure
- new symptoms occur

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs that contain acetaminophen
- 3 or more alcoholic drinks daily while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin (**Night Time only**)
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- glaucoma (**Night Time only**)
- a breathing problem such as emphysema or chronic bronchitis (**Night Time only**)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Night Time only**)

When using this product

- do not use more than directed
- excitability may occur, especially in children (**Night Time only**)
- marked drowsiness may occur (**Night Time only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Night Time only**)
- avoid alcoholic drinks (**Night Time only**)
- be careful when driving a motor vehicle or operating machinery (**Night Time only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- do not refrigerate
- store between 20-25°C (68-77°F)
- dosing cup provided

Inactive ingredients (Mucinex SINUS-MAX Day Time)

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient

Inactive ingredients (Mucinex SINUS-MAX Night Time)

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient