

Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough Caplets

Drug Facts

Active ingredients (in each caplet) Purposes

Dextromethorphan HBr 10 mg Cough suppressant
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep
 - nasal congestion due to a cold

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus) ▶

Drug Facts (continued)

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if outer package is opened or blister is torn or broken
- store at 20-25°C (68-77°F)

Inactive ingredients croscarmellose sodium, FD&C Blue #2 aluminum lake, FD&C Red #40 aluminum lake, methacrylic acid-ethyl acrylate copolymer, mica, microcrystalline cellulose, polyethylene glycol 3350, polysorbate 80, polyvinyl alcohol, povidone K29/32, sodium bicarbonate, talc, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.